

Conformal invariance of the writhe of a knot

R. Langevin

Institut de Mathématiques de Bourgogne, Université de Bourgogne

J. O'Hara*

Department of Mathematics, Tokyo Metropolitan University

March 14, 2008

Abstract

We give a new proof of the conformal invariance of the writhe of a knot from a conformal geometric viewpoint.

Key words and phrases. writhe, conformal invariant.

2000 *Mathematics Subject Classification.* Primary 57M25, Secondary 53A30.

1 Introduction

Suppose K is a framed knot, i.e. there is a unit normal vector field $\mathbf{e}_2$ along K . A 2-component link $K \cup K + \varepsilon \mathbf{e}_2$ ($|\varepsilon| \ll 1$) can be considered a closed ribbon. Let Lk and Tw be the linking number and the total twist of $K \cup K + \varepsilon \mathbf{e}_2$, and Wr the writhe of K . Then we have

$$Lk = Wr + Tw. \quad (1)$$

When the knot K has nowhere vanishing curvature and $\mathbf{e}_2$ is the principal normal vector, Lk is called the *self-linking number* of the knot and denoted by Sl . The equation (1) was proved in this case in [Că1, Că2, Că3, Po], and in [Wh] in general. It plays an important role in the application of the knot theory to molecular biology ([Ful1, Ful2, Wh-Ba]).

When the knot K is given by $K = f(S^1)$ the *writhe* is given by the Gauss integral:

$$Wr(f) = \frac{1}{4\pi} \iint_{S^1 \times S^1} \frac{\det(f'(s), f'(t), f(s) - f(t))}{|f(s) - f(t)|^3} ds dt \quad (2)$$

*This joint work was done during the second author's stay in Dijon which was supported by JSPS.

(the reader is referred to [AKT] for the details concerning writhe). Banchoff and White showed that the absolute value of the writhe is conformally invariant ([BW]). To be precise, they showed

Theorem 1.1 ([BW]) *Suppose I is an inversion in a sphere. Then we have*

$$\text{Wr}(I(K)) = -\text{Wr}(K). \quad (3)$$

It is a corollary of

Theorem 1.2 ([BW]) *Suppose I is an inversion in a sphere. Then we have*

$$\text{Tw}(I(K)) \equiv -\text{Tw}(K) \pmod{\mathbb{Z}}. \quad (4)$$

In this paper, using techniques in conformal geometry, we give new proofs of Theorem 1.1 and a special case of Theorem 1.2 when K has nowhere vanishing curvature and $\mathbf{e}_2$ is the unit principal normal vector field.

2 Notations

Throughout the paper we use the following notations.

We assume that a knot $K = f(S^1)$ is oriented. We denote the positive tangent vector $\dot{f}$ by $v = v(x)$.

We denote the circle through x, y , and z by $\Gamma(x, y, z)$. When one of x, y , and z , say z is ∞ , $\Gamma(x, y, \infty)$ means the line through x and y . When x is a point on a knot K , $\Gamma(x, x, y)$ denotes the circle (or line) which is tangent to K at x that passes through y . We assume that it is oriented by v at x . Especially, when $y = x$ $\Gamma(x, x, x)$ denotes the oriented osculating circle. (We consider lines as circles through ∞ .)

The union of osculating circles $\bigcup_{x \in K} \Gamma(x, x, x)$ is called the *curvature tube* ([BW]).

Suppose x is a point on a knot K . We denote a sphere through the osculating circle $\Gamma(x, x, x)$ and y ($y \neq x$) by $\Sigma(x, x, x, y)$. It is uniquely determined generically, i.e. unless $y \in \Gamma(x, x, x)$. When $y = \infty$ $\Sigma(x, x, x, \infty)$ means the plane through $\Gamma(x, x, x)$.

3 Proof of Theorem 1.2 in special case

In this section we assume that a knot $K = f(S^1)$ has nowhere vanishing curvature and the unit normal vector field $\mathbf{e}_2$ along K is given by the principal normal vectors. In this case the total twist is equal to $\frac{1}{2\pi}$ times the *total torsion* which we denote by $T\omega$:

$$T\omega = \frac{1}{2\pi} \int_K \tau dx,$$

where τ denotes the torsion of the knot. We will show in this section

Proposition 3.1 *If both K and $I(K)$ have nowhere vanishing curvatures then*

$$T\omega(I(K)) \equiv -T\omega(K) \pmod{\mathbb{Z}}, \quad (5)$$

where I is an inversion in a sphere.

We remark that the above proposition can be proved by Theorem 6.3 of [CSW] which gives

$$T\omega(K) = \frac{1}{2\pi} \int_K \tau dx \equiv \frac{1}{2\pi} \int_K T d\rho \pmod{\mathbb{Z}},$$

where T is the conformal torsion and ρ is the conformal arc-length.

Lemma 3.2 *Suppose a point P does not belong to the osculating circle $\Gamma(x_0, x_0, x_0)$ of a knot K at x_0 . (We allow $P = \infty$.) Then, infinitesimally speaking, the sphere $\Sigma(x, x, x, P)$ rotates around the circle $\Gamma(x_0, x_0, P)$ at $x = x_0$ as x travels in K . In other words,*

$$\lim_{x_1 \rightarrow x_0} (\Sigma(x_0, x_0, x_0, P) \cap \Sigma(x_1, x_1, x_1, P)) \supset \Gamma(x_0, x_0, P). \quad (6)$$

Proof: Let $\Sigma(x, y, z, w)$ denote a sphere through x, y, z , and w . Then

$$\begin{aligned} & \lim_{x_1 \rightarrow x_0} (\Sigma(x_0, x_0, x_0, P) \cap \Sigma(x_1, x_1, x_1, P)) \\ &= \lim_{x_1 \rightarrow x_0} \left(\lim_{y_0 \rightarrow x_0, y_1 \rightarrow x_1} (\Sigma(y_0, x_0, x_1, P) \cap \Sigma(x_0, x_1, y_1, P)) \right) \\ &\supset \lim_{x_1 \rightarrow x_0} \Gamma(x_0, x_1, P) = \Gamma(x_0, x_0, P). \end{aligned}$$

There is an alternative computational proof using the Lorentzian exterior product introduced in [LO]. $\square$

We show that the total torsion of the image of K by an inversion in a sphere with center P is equal to the total angle variation of the rotation of the sphere $\Sigma(x, x, x, P)$ as x goes around in K . In order to take into account the sign of the torsion, we have to consider the orientations.

Fix $P \notin \Gamma(x, x, x)$, where we allow $P = \infty$. The osculating circle $\Gamma(x, x, x)$ divide the sphere $\Sigma(x, x, x, P)$ into two domains. Let D_1 be one of the two that does not contain P (gray disc of Figure 1 left). Assume $\Sigma(x, x, x, P)$ is given the orientation such that the restriction to D_1 induces the same orientation to the boundary ∂D_1 as that of $\Gamma(x, x, x)$ which is fixed in the previous section. Let $n(x)$ and $n_P(x)$ be the positive unit normal vectors to $\Sigma(x, x, x, P)$ at x and P respectively (Figure 1 left). Let $\Pi(x)$ and $\Pi_P(x)$ be the normal planes to $\Gamma(x, x, P)$ at x and P respectively. We assume that $\Pi(x)$ (or $\Pi_P(x)$) is oriented so that the algebraic intersection number of $\Gamma(x, x, P)$ and $\Pi(x)$ (or $\Pi_P(x)$) at x (or respectively, at P) is equal to $+1$.

Since $\Pi(x) \perp \Gamma(x, x, P)$, Lemma 3.2 implies

Corollary 3.3 *Infinitesimally, the normal vector $n(x)$ to the sphere $\Sigma(x, x, P)$ rotates in the plane $\Pi(x)$ to K :*

$$\frac{d}{dx}n(x) \in \Pi(x). \quad (7)$$

Let I be an inversion in a sphere with center P . We denote $I(x)$ and $I(K)$ by $\tilde{x}$ and $\tilde{K}$. Then I maps the osculating circle $\Gamma(x, x, x)$ to the osculating circle $\Gamma(\tilde{x}, \tilde{x}, \tilde{x})$ of $\tilde{K}$ at $\tilde{x}$, and the sphere $\Sigma(x, x, x, P)$ to the plane $\Sigma(\tilde{x}, \tilde{x}, \tilde{x}, \infty)$. Let $\tilde{n}(\tilde{x})$ be the positive unit normal vector to $\Sigma(\tilde{x}, \tilde{x}, \tilde{x}, \infty)$. Then

$$\tilde{n}(\tilde{x}) = -n_P(x) \quad (8)$$

(Figure 1).




Figure 1:

The above convention of orientation implies that $\tilde{n}(\tilde{x})$ is equal to the unit binormal vector of $\tilde{K}$. Lemma 3.2 implies that $\tilde{n}(\tilde{x})$ rotates in $\Pi(\tilde{x})$ at $\tilde{x}$. Our convention of the orientation of $\Pi(\tilde{x})$ implies

Lemma 3.4 *The torsion of $\tilde{K}$ is equal to the angle velocity of $\tilde{n}$ with respect to the arc-length $\tilde{s}$:*

$$\tau(\tilde{K})(\tilde{x}) = \varepsilon(\tilde{x}) \left| \frac{d\tilde{n}}{d\tilde{s}}(\tilde{x}) \right| \quad (\varepsilon(\tilde{x}) \in \{+1, -1\}), \quad (9)$$

where $\varepsilon(\tilde{x})$ is the signature of the rotation of $\tilde{n}$ at $\tilde{x}$ with respect to the orientation of $\Pi(\tilde{x})$.

In other words, we have

$$\tau(\tilde{K})(\tilde{x}) = \tilde{v} \cdot \left(\tilde{n} \times \frac{d\tilde{n}}{d\tilde{s}} \right),$$

where $\tilde{v}$ denotes the positive unit tangent vector to $\tilde{K}$.




Figure 2:

Since the positive unit tangent vector to $\Gamma(x, x, P)$ at P is equal to $-\tilde{v}(\tilde{x})$, the orientations of $\Pi_P(x)$ is opposite to that of $\Pi(\tilde{x})$. Therefore, (8) and Lemma 3.4 imply

$$\tau(\tilde{K})(\tilde{x}) = -\varepsilon_P \left| \frac{ds}{d\tilde{s}} \right| \left| \frac{dn_P}{ds} \right| \quad (\varepsilon_P \in \{+1, -1\}),$$

where ε_P is the signature of the rotation of n_P with respect to the orientation of $\Pi_P(x)$. Then Lemma 3.2 implies $\left| \frac{dn_P}{ds} \right| = \left| \frac{dn}{ds} \right|$ (Figure 3), therefore we have

Lemma 3.5 *The torsion of $\tilde{K}$ at $\tilde{x}$ is equal to the negative of the angle velocity of n in $\Pi(x)$ with respect to the arc-length s of K up to the multiplication by the Jacobian:*

$$\tau(\tilde{K})(\tilde{x}) = -\varepsilon(x) \left| \frac{ds}{d\tilde{s}} \right| \left| \frac{dn}{ds}(x) \right| \quad (\varepsilon(x) \in \{+1, -1\}), \quad (10)$$

where $\varepsilon(x)$ is the signature of the rotation of n at x with respect to the orientation of $\Pi(x)$.

In other words, we have

$$\tau(\tilde{K})(\tilde{x}) = - \left| \frac{ds}{d\tilde{s}} \right| v \cdot \left(n \times \frac{dn}{ds} \right).$$

Thus we are led to




Figure 3: $\angle n(x) \cdot n(x_1) = \angle n_P(x) \cdot n_P(x_1) = \angle \Sigma(x, x, x, P) \cdot \Sigma(x_1, x_1, x_1, P)$

Proposition 3.6 *Let K be a knot with nowhere vanishing curvature, I an inversion in a sphere with center P , and $\tilde{K} = I(K)$. Assume that $\tilde{K}$ has nowhere vanishing curvature, which happens if and only if P is not contained in the curvature tube of K . Then the total torsion of $\tilde{K}$ is the negative of the total angle variation of the positive unit normal vector $n(x)$ to the sphere $\Sigma(x, x, x, P)$ as x goes around in K :*

$$\int_{\tilde{K}} \tau(\tilde{K})(\tilde{x}) d\tilde{x} = - \int_K \varepsilon |dn| = - \int_K v \cdot (n \times dn). \quad (11)$$

Corollary 3.7 *Let K be a knot with nowhere vanishing curvature, and I_j ($j = 1, 2$) an inversion in a sphere with center P_j which is not contained in the curvature tube of K . Then the total torsion of $I_1(K)$ and $I_2(K)$ coincide modulo $2\pi\mathbb{Z}$:*

$$\int_{I_1(K)} \tau d\tilde{x}_1 \equiv \int_{I_2(K)} \tau d\tilde{x}_2 \pmod{2\pi\mathbb{Z}}.$$

In other words,

$$T\omega(I_1(K)) - T\omega(I_2(K)) \in \mathbb{Z}.$$

Proof: Let n_j ($j = 1, 2$) be the positive unit normal vector to $\Sigma(x, x, x, P_j)$ at x , and θ_{21} the angle from n_1 to n_2 in the oriented plane $\Pi(x)$. Then Corollary 3.3 and the above Proposition imply

$$\int_{I_1(K)} \tau d\tilde{x}_1 - \int_{I_2(K)} \tau d\tilde{x}_2 = \int_K d\theta_{21},$$

which is equal to $2\pi k$ ($k \in \mathbb{Z}$) since K is closed. □

Proof of Proposition 3.1: Put $P_2 = \infty$ in the above Corollary. □

4 Proof of Theorem 1.1

We first prove the following fact in our context, which implies Theorem 3.1 in the case when the knot has nowhere vanishing curvature.

Proposition 4.1 ([Că1], [Po]) *Suppose K has nowhere vanishing curvature. Then*

$$4\pi \text{Wr}(K) = -2 \int_K \tau + 4\pi k \quad \text{for some } k \in \mathbb{Z}.$$

In other words, $\text{Wr} + T\omega$ is an integer.

Proof: Assume $K = f(S^1)$ is parametrized by the arc-length. Suppose $S^1 = [0, L]/\sim$, where L is the length of K . Let φ be a map from $S^1 \times S^1 \setminus \Delta$ to S^2 given by

$$\varphi(s, t) = \frac{f(s) - f(t)}{|f(s) - f(t)|} \quad (s \neq t).$$

Then the integrand of the writhe is equal to the pull-back of the standard area element of S^2 by φ :

$$\frac{\det(f'(s), f'(t), f(s) - f(t))}{|f(s) - f(t)|^3} = \varphi \cdot (\varphi_s \times \varphi_t),$$

where $\varphi_s = \frac{\partial \varphi}{\partial s}$ and $\varphi_t = \frac{\partial \varphi}{\partial t}$. Therefore, 4π times the writhe is equal to the signed area of the image of

$$D = \{(s, t) \mid 0 \leq s \leq L, s < t < s + L\}$$

by φ . Let S be the closure of $\varphi(D)$. Then S is an oriented “surface” (a continuous image of $S^1 \times [0, L]$) possibly with self-overlaps in S^2 whose “boundary” (the image of $S^1 \times \{0, L\}$) is given by

$$\partial S = C_+ \cup (-C_-),$$

where C_+ and C_- denote the positive and negative tangential indicatrices:

$$C_+ = \varphi(\Delta_+) = \{v(s) = \dot{f}(s) \mid 0 \leq s \leq L\},$$

$$C_- = \varphi(\Delta_-) = \{-v(s) = -\dot{f}(s) \mid 0 \leq s \leq L\}.$$

We assume that the indicatrices are oriented so that s increases in the positive direction.

Let $\Gamma(s)$ denote a great circle in S^2 which is tangent to C_+ at $v(s)$ (and to C_- at $-v(s)$). We remark that C_+ has nowhere vanishing tangent vector since the knot has nowhere vanishing curvature. Note that $C_+ \cup C_-$ is the envelope of the family $\{\Gamma(s)\}_{0 \leq s \leq L}$. We assume that $\Gamma(s)$ has an orientation compatible with that of C_+ at $v(s)$. Let $\Gamma_+(s)$ be a semi-circle of $\Gamma(s)$ from $v(s) \in C_+$ to

$-v(s) \in C_-$ in the positive direction of $\Gamma(s)$, and S' a region in S^2 swept by $\Gamma_+(s)$ as s varies in $[0, L]$. Then S' is given by

$$S' = \{w(s, t) = (\cos t)v(s) + (\sin t)\frac{\dot{v}}{|\dot{v}|}(s) \mid 0 \leq s \leq L, 0 \leq t \leq \pi\}.$$

It is an oriented “surface” (a continuous image of $S^1 \times [0, \pi]$) possibly with self-overlaps in S^2 whose “boundary” (the image of $S^1 \times \{0, \pi\}$) is $C_+ \cup (-C_-)$. The signed area of S' is given by

$$\text{Area}(S') = \int_0^L \int_0^\pi w \cdot (w_s \times w_t) ds dt.$$

Since $\dot{v}/|\dot{v}|$ is equal to the principal normal vector e_2 of the knot K , we have

$$\begin{aligned} w \cdot (w_s \times w_t) &= \det((\cos t)e_1 + (\sin t)e_2, (\cos t)\kappa e_2 + (\sin t)\dot{e}_2, -(\sin t)e_1 + (\cos t)e_2) \\ &= -\tau \sin t, \end{aligned}$$

which implies

$$\text{Area}(S') = -2 \int_0^L \tau ds = -4\pi T\omega(K).$$

Since both S and S' have the boundary $C_+ \cup (-C_-)$, $S \cup (-S')$ is a cycle of S^2 , i.e. an oriented “surface” (a continuous image of a torus) possibly with self-overlaps without a boundary. Therefore, the signed area of $S \cup (-S')$ is equal to $4\pi k$ for some integer k . It follows that

$$\text{Area}(S \cup (-S')) = \text{Area}(S) - \text{Area}(S') = 4\pi Wr + 4\pi T\omega = 4\pi k \quad (k \in \mathbb{Z}),$$

which completes the proof. $\square$

Corollary 4.2 *If both K and $I(K)$ have nowhere vanishing curvatures then*

$$Wr(I(K)) = -Wr(K).$$

Proof: Let I_j ($j = 0, 1$) be an inversion in a sphere with center P_j which is not contained in the curvature tube of K . Then Proposition 4.1 implies that

$$\begin{aligned} Wr(I_0(K)) + T\omega(I_0(K)) &\in \mathbb{Z}, \\ Wr(I_1(K)) + T\omega(I_1(K)) &\in \mathbb{Z}. \end{aligned}$$

Then Corollary 3.7 implies

$$Wr(I_0(K)) - Wr(I_1(K)) \in \mathbb{Z}.$$

Join P_0 and P_1 by a smooth path P_t . Let I_t be an inversion in a sphere with center P_t . Then $Wr(I_t(K))$ is a continuous function of t ($[Po]$), and hence

$$Wr(I_0(K)) = Wr(I_1(K)).$$

When P_1 goes to ∞ and the radius of the sphere of the inversion also goes to $+\infty$, $I_1(K)$ approaches the mirror image of K and hence $\text{Wr}(I_1(K))$ approaches $-\text{Wr}(K)$, which completes the proof. $\square$

Proof of Theorem 1.1: Suppose the curvature of K vanishes somewhere. We have only to show that K can be approximated, with respect to the C^2 -topology, by a knot with non-vanishing curvature. This can be done as follows. The curvature tube of K is non-compact as it contains a line. But we can still find a point P with a very big distance from K which is not contained in the curvature tube. Let I be an inversion in a sphere with center P and radius approximately equal to the distance between P and K . We can get a desired knot by taking the mirror image of $I(K)$ thus constructed. $\square$

References

- [AKT] J. Aldinger, I. Klapper, and M. Tabor, *Formulae for the calculation and estimation of writhe*, J. Knot Theory Ramifications. **4** (1995), 343–372.
- [BW] T. Banchoff and J. H. White, *The behavior of the total twist and self-linking number of a closed space curve under inversions*, Math. Scand. **36** (1975), 254–262.
- [Că1] G. Călugăreanu, *L'intégrale de Gauss et l'analyse des nœuds tridimensionnels*, Rev. Math. Pures Appl. **4** (1959), 5–20.
- [Că2] G. Călugăreanu, *Sur les classes d'isotopie des noeuds tridimensionnels et leur invariants*, Czechoslovak Math. J. **11** (1961), 588–625.
- [Că3] G. Călugăreanu, *O teoremă asupra înlănțuirilor tridimensionale de curbe închise*, Comm. Acad. R. P. Romîne (1961), 829–832.
- [CSW] G. Cairns, R. W. Sharpe and L. Webb, *Conformal invariants for curves and surfaces in three dimensional space forms*, Rocky Mountain Journal of Math. **24** (1994), 933–959.
- [Ful1] F. B. Fuller, *The writhing number of a space curve*, Proc. Natl. Acad. Sci. USA **68** (1971), 815–819.
- [Ful2] F. B. Fuller, *Decomposition of the linking of a closed ribbon: a problem from molecular biology*, Proc. Natl. Acad. Sci. USA **75** (1978), 3557–3561.
- [LO] R. Langevin and J. O'Hara, *Conformally invariant energies of knots*, J. Inst. Math. Jussieu **4** (2005), 219–280.
- [Po] W. F. Pohl, *The self-linking number of a closed space curves*, J. of Math. Mech. **17** (1968), 975–985.
- [Wh] J. H. White, *Self-linking and the Gauss integral in higher dimensions*, Amer. J. Math. **91** (1969), 693–728.

[Wh-Ba] J. H. White and W. R. Bauer, *Calculation of the twist and the writhe for representative models of DNA*, J. Mol. Biol. **189** (1989), 329–341.